

# SHARP Health Plan

Men (18 years and older)<sup>1</sup>

## Preventive Care Checklist

No matter how healthy you are today, having a discussion with your doctor about preventive care can make a difference in the long run — for you and the people you care about. Discuss your checklist with your doctor to develop your own personalized preventive care plan. Your doctor will help you determine which tests and health screenings are best for you based on your age, gender, health status and family history.<sup>1</sup>

Visits		
Welcome to Medicare preventive visit (once during first 12 months with Part B)	18 and older	<input type="radio"/>
Annual Wellness Visit (does not include physical exam)	18 and older	<input type="radio"/>
Vaccines		
Flu (each flu season)	18 and older	<input type="radio"/>
Hepatitis B (those with medium to high risk)	18 and older <sup>2</sup>	<input type="radio"/>
Pneumococcal	over 65 or younger with risk factors	<input type="radio"/>
COVID-19	18 and older	<input type="radio"/>
Shingrix series (once)	50 and older <sup>3</sup>	<input type="radio"/>
Diphtheria, tetanus and pertussis (every 10 years)	18 and older <sup>3</sup>	<input type="radio"/>
Screenings		
Abdominal aortic screening ultrasound	18 and older <sup>4</sup>	<input type="radio"/>
Colorectal cancer screening, may include one or more of the following: <ul style="list-style-type: none"> <li>• Fecal occult blood tests (once every 12 months)</li> <li>• Flexible sigmoidoscopies (once every 48 months, or 120 months after previous screening colonoscopy)</li> <li>• Colonoscopy (once every 120 months, or every 24 months if high risk, or 48 months after screening flexible sigmoidoscopy)</li> <li>• Barium enemas (once every 48 months)</li> <li>• Multi-target stool DNA test / Cologuard (once every 3 years with no symptoms and average risk)</li> <li>• Blood-based markers, if available (once every 3 years with no symptoms and average risk)</li> </ul>	50 and older	<input type="radio"/>

## Men's preventive care checklist, continued<sup>1</sup>

Screenings		
Diabetes screening (2 per year with risk factors)	18 and older with risk factors	<input type="radio"/>
Glaucoma screening (every 12 months)	18 and older with risk factors	<input type="radio"/>
HIV screening (every 12 months) if ages	15-65 or at high risk	<input type="radio"/>
Lung cancer screening with low-dose CT (once yearly)	55-77 <sup>2</sup>	<input type="radio"/>
PSA blood test (once a year)	50 and older	<input type="radio"/>
Bone mass measurements (every 24 months)	50-85 with certain risk factors	<input type="radio"/>
Alcohol misuse counseling (PCP)	18 and older	<input type="radio"/>
Cardiovascular behavioral therapy (PCP) includes blood pressure, diet advice and discussion to lower risk for heart attacks and stroke	18 and older	<input type="radio"/>
Cardiovascular disease screening (once every 5 years, including cholesterol, lipid and triglyceride levels)	15-65 or at high risk	<input type="radio"/>
Tobacco prevention and counseling	18 and older	<input type="radio"/>
Depression screening yearly (PCP)	18 and older	<input type="radio"/>
Hepatitis B screening tests if high risk (PCP)	18 and older	<input type="radio"/>
Hepatitis C screening tests (yearly if high risk, one time if born 1945-1965 or if you had a blood transfusion before 1992)	18 and older	<input type="radio"/>
Obesity behavioral therapy (PCP)	18 and older	<input type="radio"/>
Sexually transmitted infection screen and counseling (PCP)	18 and older	<input type="radio"/>

<sup>1</sup> The information in this guide includes recommendations adapted from the following sources as of November 2022 and is subject to change: preventive services with a rating of A or B from the U.S. Preventive Services Task Force; immunization for children, adolescents and adults recommended by the Centers for Disease Control and Prevention; and preventive care screenings for infants, children, adolescents and women supported by the Health Resources and Services Administration. <sup>2</sup> Based on risk factors, and as advised by your doctor. <sup>3</sup> This is a Part D vaccine with \$0 copay. <sup>4</sup> Once if family history of abdominal aortic aneurysm, or for males ages 65-75 and who have smoked at least 100 cigarettes in their lifetime.